First Hit

Previous Doc

Next Doc

Go to Doc#

Generate Collection Print

L20: Entry 43 of 45

File: JPAB

Jun 20, 2000

PUB-NO: JP02000169502A

DOCUMENT-IDENTIFIER: JP 2000169502 A

TITLE: BRANCHED MALTODEXTRIN AND ITS PRODUCTION

PUBN-DATE: June 20, 2000

INVENTOR-INFORMATION:

NAME

POURCUE CAMUEDINE

COUNTRY

FOUACHE, CATHERINE DUFLOT, PIERRICK LOOTEN, PHILIPPE

ASSIGNEE-INFORMATION:

NAME

COUNTRY

ROQUETTE FRERES

APPL-NO: JP11345503

APPL-DATE: December 3, 1999

PRIORITY-DATA: 1998FR-15344 (December 4, 1998)

INT-CL (IPC):  $\underline{C08} \ \underline{B} \ \underline{30/18}; \ \underline{A23} \ \underline{G} \ \underline{3/30}$ 

ABSTRACT:

PROBLEM TO BE SOLVED: To provide a new-type branched <u>maltodextrin</u> specified in a 1,6-glucoside bond level, a reducing sugar content, a polydispersity index, and an average molecular weight and being usable as a product intended for <u>digestion</u> by man or animals.

SOLUTION: Provided are a branched maltodextrin and a hydrogenated maltodextrin each having a 1,6-glucoside bond level of 22-35%, a reducing sugar content of at most 20%, a polydispersity index of at most 5, and a number- average molecular weight equal to at most 4,500 g/mol. This (hydrogenated) branched maltodextrin is obtained by treating an acidified starch dehydrated so as to have a water content of at most 5% at 120-300°C in a thin-layer continuous reactor, collecting, purifying, and, desirably, concentrating the branched derived starch product, fractionating the product according to numberaverage molecular weights, and, optionally, catalytically hydrogenating the obtained product. This in combination with a polyol is utilized as a non-cariogenic composition. In use, 30-70 wt.% branched maltodextrin is mixed with 30-70 wt.% multitol.

COPYRIGHT: (C) 2000, JPO

Previous Doc Next Doc Go to Doc#